

Review

Signaling via NF- κ B in the nervous system

Barbara Kaltschmidt, Darius Widera, Christian Kaltschmidt*

Institut für Neurobiochemie Universität Witten/Herdecke, Stockumer Street 10, D-58448 Witten, Germany

Received 9 March 2005; received in revised form 20 May 2005; accepted 23 May 2005

Available online 16 June 2005

Abstract

Nuclear factor kappa B (NF- κ B) is an inducible transcription factor present in neurons and glia. Recent genetic models identified a role for NF- κ B in neuroprotection against various neurotoxins. Furthermore, genetic evidence for a role in learning and memory is now emerging. This review highlights our current understanding of neuronal NF- κ B in response to synaptic transmission and summarizes potential physiological functions of NF- κ B in the nervous system. This article contains a listing of NF- κ B activators and inhibitors in the nervous system, furthermore specific target genes are discussed. Synaptic NF- κ B activated by glutamate and Ca^{2+} will be presented in the context of retrograde signaling. A controversial role of NF- κ B in neurodegenerative diseases will be discussed. A model is proposed explaining this paradox as deregulated physiological NF- κ B activity, where novel results are integrated, showing that p65 could be turned from an activator to a repressor of anti-apoptotic genes.

© 2005 Elsevier B.V. All rights reserved.

Keywords: Transcription factor; Neuron; Neuroprotection; Memory; Tumor necrosis factor; Calcium; Synapse; Retrograde; Inhibitor; Activator; Target gene; NF- κ B; I κ B

1. NF- κ B activation via tumor necrosis factor (TNF)

Nuclear factor kappa B (NF- κ B) was discovered by David Baltimore's laboratory as an inducible transcription factor in lymphocytes [1]. The so-called canonical pathway of NF- κ B activation via tumor necrosis factor (TNF) [2,3] will be presented to summarize part of the general knowledge on activation mechanisms (see Fig. 1). The TNF pathway is one of the best characterized NF- κ B-dependent pathways with more than 70 000 references in PubMed. Therefore, we have chosen this so-called canonical pathway, which is also operating in the nervous system, to introduce the NF- κ B activation.

Within the nervous system, TNF (a 17 kDa protein) can bind to TNF receptors (TNF-Rs) expressed on both glia and neurons [4]. The expression of the TNF- α gene is subject to autoregulation via activated NF- κ B [5]. Two different receptors, p55 (TNF-R1) and 75 (TNF-R2), have been identified. The p55 receptor is thought to be the major NF- κ B activating TNF-R [6].

Central to NF- κ B activation seems to be the I κ B kinase complex (IKK), which is catalyzing the signal-dependent phosphorylation of the NF- κ B inhibiting I κ B. Thus, the IKK complex initiates NF- κ B activation via phosphorylation of I κ B, which is a signal for polyubiquitination and subsequent degradation. An amazing wealth of information has been accumulated, describing how receptor activation might be connected to IKK activation. There are themes of ubiquitination after the activation of the trimeric TNF receptor. The binding of soluble or cell-bound TNF leads to the activation of the latent trimerized TNF receptors. These receptors share an intracellular so-called death domain (DD) with several other TNF receptors such as TRAIL receptors or DR3, DR6 and various other receptors with non-TNF ligands such as the CD 95 (Apo/Fas) receptor or the p75 low-affinity nerve growth factor receptor. TNF-R1 is unique in its composition of intracellular interaction proteins [7]. The genetic ablation of TNF-R1 (p55) exacerbates traumatic brain injury and correlated with a reduced NF- κ B activation [4]. Recent data suggest that the TNF-R2 is responsible for a persistent NF- κ B activation and neuroprotection [8,9].

* Corresponding author.

E-mail address: c.kaltschmidt@uni-wh.de (C. Kaltschmidt).




Fig. 1. Canonical pathway of NF- κ B activation by tumor necrosis factor (TNF). The activation of the TNF receptor by ligand binding is transmitted to the IKK complex which phosphorylates I κ B family inhibitory molecules (see text for details). This targets I κ B for polyubiquitination and subsequent degradation within the 26S proteasome. This frees nuclear localization signal on the DNA-binding p65/p50 subunits. After nuclear import, target gene transcription is initiated. Not all signaling components depicted in the canonical pathway have been investigated in the nervous system, but appear to be present in neurons and glia. Proteins are depicted as icons which illustrate a functional category (receptor, enzyme, etc.) as suggested by the Alliance for Signalling convention (www.signaling-gateway.org). Ub=ubiquitination; P=phosphorylation.

The trimerization of the non-signaling TNF-R complex is mediated by a N-terminal pre-ligand assembly domain. TNF binding activates the pre-assembled receptor trimer via release of silencer of death domains (SODD) from the intracellular death domains; see [7] for discussion. The physiological role of SODD remains unclear. The trimeric death domains seem to function as an assembly platform for further intracellular interactors such as the adapter protein TRADD. TRADD seems to enable a bifurcation in either the apoptotic pathway, leading to caspase activation, or in the anti-apoptotic NF- κ B-dependent pathway, which involves the transcription of genes encoding survival factors. Apoptosis, as well as anti-apoptosis, is involving TRADD in the context of different signaling complexes [10]. The initial receptor-bound complex (see Fig. 1) might contain TRADD, RIP and TRAF2. This provides a signaling scaffold for the activation of NF- κ B. An apoptotic pathway could be triggered by FADD, which might interact with RIP and

TRADD in a non-receptor bound cytoplasmic complex. With its N-terminal domain, TRADD could facilitate the interactions with TRAF1 and TRAF2. Anti-apoptotic proteins are targeted to the receptor complex, e.g. cIAP-1 and cIAP-2. This might be the reason for the relative good protection of activated TNF-R1 signaling against apoptosis. Several studies with murine neuronal cultures use human TNF- α , which has been shown to activate the TNF-R1 [11,12].

The I κ B kinase complex might integrate many NF- κ B activating stimuli culminating on NF- κ B activation within the nervous system such as TNF, LPS, IL-1, NGF or glutamate-mediated signaling; see [13] for a review.

Crucial for NF- κ B activation is the phosphorylation of I κ B. This is most commonly due to the interaction of activated I κ B-kinase complex (IKK) with I κ B. The IKK complex is composed of two catalytic subunits (IKK- α and IKK- β), a receptor targeting/oligomerization subunit NEMO and ELKS, a recently discovered I κ B- α targeting subunit [14]. The activation of the IKK complex might be regulated by multiple mechanisms [2]. A classical phosphorylation of an activation loop has been reported for the main I κ B phosphorylating kinase IKK- β at serines 177 and 181 [15]. This might be either due to autophosphorylation or due to upstream kinases. Surprisingly, only for the upstream kinase MEKK3 genetic evidence for an involvement in the TNF pathway could be provided [16]. Other activation mechanisms dependent on the oligomerization of the IKK complex have been described [17,18]. NEMO-mediated recruitment of the IKK complex to the T cell receptor complex has been identified as activation mechanism [19]. Surprisingly, also the membrane localization of NEMO could activate the IKK complex [19]. NEMO, the regulatory subunit of the IKK complex, is inducibly ubiquitinated. The receptor-bound complex containing TRAF2 and TRAF5 could recruit the IKK complex to the membrane-bound TNF receptor and thus lead to activation via IKK oligomerization [20]. I κ B is recruited to the IKK complex via an interaction with ELKS [14]. The activated IKK complex could catalyze the phosphorylation of Ser 32 and 36 on the I κ B- α molecule.

As summarized above, the activation of the IKK complex seems to rely on multimerization. The ubiquitination of NEMO might be an essential prerequisite of the activation process, since deubiquitinating enzymes such as CYLD [21,22] are essential to deactivate the IKK complex. These ubiquitination pathways seem to be independent of the proteasomal degradation but are modulating oligomeric state. Other deubiquitination and ubiquitination activity is found to be encoded in the protein A20, which could target RIP for degradation [23].

For an extensive review of the ubiquitin-proteasome system, see the special issue of BBA, Vol. 1695 (2004). Ubiquitin (Ub) is a small (86 kDa) protein used to tag proteins either for the degradation or for signaling (multimerization). Ubiquitin is conjugated to the amino groups of lysine residues on target proteins by a cascade of enzymes called E1, E2 and E3 [24]. An SCF (Skp-1/Cul/F box)-type

multisubunit E3 ubiquitin ligase holoenzyme contains the phospho-I κ B specific acceptor subunit bTrCF and is responsible for I κ B poly-ubiquitination [25]. Interestingly, one of the frequently used pharmacological inhibitors of NF- κ B activation pyrrolidone dithiocarbamate (PDTC) acts as an inhibitor of the I κ B ubiquitin–ligase complex [26]. The signal for the ubiquitin–ligase seems to be the phosphorylation of serines 32 and 36 on I κ B- α .

I κ B proteins are essential regulators of nuclear import, which can interact with the nuclear localization signal (NLS) to induce an alpha helical conformation [27,28]. Interestingly, I κ B- α only interacts with the NLS of p65, whereas I κ B- β interacts with both the NLS of p50 and p65 [29]. This conformation cannot be recognized by the nuclear import receptor importin alpha [27], in contrast, I κ B free nuclear localization signal assumes a random coil conformation, which is the basis for the interaction with importin. Recently, it was shown that the NLS of p50 and p65 were recognized mainly by importin α 3 [30]. A fresh view on import/export suggests additional complexity since the trimeric p50/p65 I κ B- α complex is shuttled between nucleus and cytoplasm [29]. The nucleocytoplasmic shuttling of the trimeric complex of p50, p65 and I κ B- α could not be observed by [30]. In addition to inhibition by I κ B, there seems to be a novel mechanism to terminate NF- κ B signaling: promoter-specific degradation of p65 via nuclear proteasomes [31].

2. NF- κ B subunits

Inducible NF- κ B resides in the cytoplasm in a latent DNA-binding form (see Fig. 1: blue p50 and p65 molecules). So far, five DNA-binding isoforms have been described: RelB, c-Rel, p65 (RelA), p50 and p52, the latter two being proteolytic products of larger precursors. Surprisingly, p50 and p52 are products of specific protein degradation within the proteasome. In some proteins, the proteasome only degrades specific parts of the protein and leaves the rest intact. This process, named regulated ubiquitin proteasome-dependent pathway (RUP), was recently reviewed by [32]. The C-terminal part of the NF- κ B p50 precursor p105 is degraded by the proteasome in RUP-dependent manner. Degradation occurs either co- or posttranslationally. Mechanistically, a protection of degradation via dimerization has been suggested. Interestingly, the dimerization of NF- κ B can already occur between two nascent polypeptide chains emerging from successive ribosomes on the mRNA [33]. This complex way of processing might add further ways of regulation, especially in neurons where proteasome activity seems to be regulated by synaptic activity [34]. The NF- κ B subunits RelB, c-Rel and p65 (RelA) contain transactivating domains that are capable to activate transcription without the help of other NF- κ B subunits. DNA-binding occurs as a dimer, whereas the latent non-DNA-binding NF- κ B complex contains an

inhibitory subunit called I κ B. Later on, more I κ B subunits were described. Well-characterized inhibitory subunits are: I κ B- α , I κ B- β , I κ B- γ (p105), I κ B- δ (p100) and I κ B- ϵ [35].

3. NF- κ B within the nervous system

Central functions of the nervous system are information transmission, processing and storage. To fulfill these elaborated demands, special cellular structures have been evolved. Neurons are able to receive, transmit and store information. Two neurons can communicate via synapses, which are specialized cellular compartments, consisting of a presynaptic (sending) cell and a postsynaptic (receiving) cell. Frequently, chemical synapses (see Fig. 2) are used to communicate within the neuronal networks. Other important cell types are summarized as glia (neuronal glue). These cells can assist or participate in synaptic information processing. Neurodegeneration can destroy subcellular structures such as synapses and/or lead to neuronal cell death. Neuroprotective processes include calcium buffering, generation of novel synapses, anti-apoptotic gene expression, caspase inhibition, balancing of reactive oxygen intermediates, etc., thus protecting neurons from neurodegeneration.




Fig. 2. Localization of the NF- κ B activation machinery within the synapse. Both pre- and postsynapse seem to contain the necessary components. The importance of pre- and postsynaptic activation mechanisms has to be clarified.

Table 1
Examples for molecules activating NF- κ B in the nervous system

Molecule	Cell type	Reference
Glutamate	Neurons	[42]
Kainate	Neurons	[41]
NMDA	Neurons	[52]
TNF	Neuroblastoma	[53]
TNF	Neurons	[54,55]
TNF	Astrocytes	[56]
TNF	Microglia	[57]
IL-1	Glia cells	[13]
IL-1	Neurons	[58]
sAPP	Neurons	[59]
sAPP	Microglia	[60]
A β	Neurons	[43,61]
A β	Astrocytes	[43]
A β /IFN γ	Microglia	[62]
ATP	Microglia	[63]
ATP/IL-1	Astrocytes	[64]
Adenosine	Neurons	[65]
LPS	Microglia	[66]
LPS	Astrocytes	[67]
EPO	Neurons	[68]
Bradykinin	Neurons	[69]
SDF-1	Astrocytes	[70]
EGF	Astrocytes	[71]
VEGF	Neuroblastoma	[72]
NGF	Neurons	[73]
NGF	Schwann C.	[49]
NGF	Oligod.	[74]
CNTF	PNS neurons	[75]
LIF	PNS neurons	[75]
IL-6	PNS neurons	[75]
CT-1	PNS neurons	[75]
BDNF	Microglia	[50]
NT-3	Microglia	[50]
NT4/5	Microglia	[50]
IGF-1	Neuroblastoma	[76]
ADNF	Neuroblastoma	[77]
PEDF	Neurons	[78]
H ₂ O ₂	Neurons	[55]
H ₂ O ₂	Oligodendrocyte	[79]
Gangliosides	Neurons	[80]
Glutaredoxin	Neurons	[81]
Focal cerebral ischemia	Neurons	[82]
Injury	Neurons	[130]
RAGE dependent diabetic neuropathy	Neurons	[83]
Non-A β amyloid	Neurons	[84]
Amitriptyline	Neurons	[85]
Desipramine	Neurons	[85]
Fluoxetine	Neurons	[85]
IL-1	Neurons	[86]
Lack of prion protein	Neurons	[87]
Selenium deficiency	Neurons	[88]
Intracellular calcium	Neurons	[89]
AMPA	Neurons	[90]
BDNF	Neurons	[91]
NGF	Neurons	[91]
NT-3	Neurons	[91]
NT-4/5	Neurons	[91]
Pre-myelination	Schwann cells	[92]
Dehydroepiandrosterone (DHEA)	PC-12	[93]
Allopregnanolone (Allo)	PC-12	[93]
Sleep deprivation	Neurons	[94]
FAIM	Neurons	[95]
TGF- β 1	Neurons	[96]

Within the nervous system, inducible NF- κ B is most frequently composed of 2 DNA-binding subunits (e.g., p50 or p65) that are either constitutively active or form a complex with the inhibitory subunit I κ B- α [36–43]. There are reports of other κ B-binding activities such as brain-specific transcription factor (BETA) specifically detected in grey matter extracts [44], developing brain factors (DBFs), which were reported to be highly enriched in developing cortex [45] and neuronal κ B binding factor (NKBF) with different target sequence requirements [46]. These κ B binding factors were not assigned to specific genes, nor could they be tested directly in reporter gene assays. On the other hand, it seems to be granted that there is an additional level of complexity, added by overlapping mutually exclusive or synergistically acting binding sites for other transcription factors. In glia and neurons, a sustained NF- κ B activation for up to 72 h was observed. Novel data suggest that this might be due to a differential use of I κ B isoforms α and β . For glial cells, it was reported that sustained NF- κ B activity induced via IL-1 was still present even after I κ B- α protein amount was returned to the level measured before stimulation, whereas I κ B- β protein levels remained low, suggesting that I κ B- β is the negative regulator for sustained NF- κ B activation [47]. In contrast, a biphasic response which is repressed by I κ B- α was reported for TNF-stimulated neural cells [48]. Whereas a lot of knowledge on the mechanisms of TNF-mediated signaling has been accumulated from non-neuronal cells, a systematic analysis of the TNF pathways within the nervous system has not been done so far.

4. Activators of NF- κ B within the nervous system

Many activators of NF- κ B were identified (Table 1), some with specificity for the nervous system such as the neurotransmitter glutamate acting as an NF- κ B activator via the main ionotropic glutamate receptors or the neurotrophins, which display an amazing specificity: NGF is activating NF- κ B via the p75 receptor [49], whereas other neurotrophins such as NT-3 or NT4/5 do not activate NF- κ B. In microglia, all neurotrophins do activate NF- κ B [50]. Taken together, there is a lot of complexity in the system: One molecule like TNF can activate or repress NF- κ B in neurons [51], and there are cell-type specific effects.

5. Inhibitors of NF- κ B within the nervous system

There are anti-inflammatory cytokines which are known from the immune system like TGF β or IL-10 which also

Notes to Table 1:

NMDA: *N*-methyl-D-aspartate; TNF: tumor necrosis factor; IL-1: interleukin-1; sAPP: secreted beta-amyloid precursor; A β : beta-amyloid; IFN γ : interferon γ ; LPS: lipopolysaccharide; EPO: erythropoietin; SDF-1: stromal-derived cell factor-1 α ; EGF: epidermal growth factor; AMPA: alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid.

inhibit NF- κ B in the nervous system (see Table 2). Mechanistically, it is less clear how these molecules act in the nervous system. One possibility might be the induction of I κ B transcription [97]. Some molecules are already known as activators, but seem to act in higher concentration as repressors. Interestingly, the lipid peroxidation product 4-hydroxy-2,3-nonenal inhibits both constitutive and inducible NF- κ B activity [98].

6. Target genes within the nervous system

In a heroic effort, H.L. Pahl compiled 150 stimuli and 150 target genes of NF- κ B [123]. This ever growing list of target genes, mainly in non-neuronal cells, is updated on T.D. Gilmore's webpage (www.nf-kb.org). A limited number of genes regulated by brain NF- κ B and with direct relevance for the nervous system have so far been described [124]. These include the neural cell adhesion molecules [125], inducible nitric oxide synthase [126], amyloid precursor protein [127], μ -opioid receptors [128], brain-derived neurotrophic factor [52], inducible cyclooxygenase 2 (Cox II, [129]) and Ca²⁺/calmodulin-dependent protein kinase II (CamKII δ [130]).

Table 2
Examples for molecules repressing NF- κ B in the nervous system

Molecule	Cell type	Reference
IL-4	Astrocytes	[99]
IL-10	Astrocytes	[99]
IL-10	Neurons	[100]
IL-10	Microglia	[101]
TGF β	Microglia	[102]
TGF β	Neurons	[103]
TNF/H ₂ O ₂	Neurons	[104]
TNF	Neurons	[55]
NO	Neurons	[105]
NO	Microglia	[106]
Hydroxy-nonenal	Neurons	[98]
Glucocorticoids	Neurons	[107]
A β	Neurons	[43]
Melatonin	Neurons	[108]
Corticotropin-releasing-hormone	Neurons	[109]
Aspirin	Neurons	[58]
Triflusal	Glia	[110]
LY341122	Neurons	[111]
Ganglioside	Astrocyte	[112]
PDTC	Neurons	[41]
PDTC	Microglia	[66]
Vit E	Neurons	[113]
Dexanabinol (cannabinoid)	Neurons	[114]
Hypericin	Neurons	[115]
Selegiline	Neuroblastoma	[116]
Methylpyridinium (MPP(+))	Neuroblastoma	[117]
Silymarin	Microglia	[118]
Mutant preseniline-1	Neurons	[119]
Caffeic acid phenethyl ester (CAPE)	Neurons	[120]
Selenite	Neurons	[121]
BAY 11 7082	Neurons	[122]

To our knowledge, there are only few studies analyzing NF- κ B-dependent gene expression in the nervous system on larger scales. In a microarray study using U373 human glioblastoma cells, many genes responded to TNF (>880 from 7500 tested) and were measured with a more than two-fold induction rate [131]. In this study, several novel TNF-responsive genes (about 60% of the genes regulated by a factor >3) were detected. A comparison of TNF-induced gene expression profiles of nervous system derived U373 cells, with profiles from non-nervous system 3T3 and Hela cells revealed striking cell-type specificity. Several of the TNF-induced genes were repressed by the inhibitor of IKK: PDTC and might therefore constitute novel target genes. It is not easy to discriminate between a role for NF- κ B in glia or neurons in vivo. Cell-type specific knockout models might help to clarify this issue. Recently, a study using brain ischemia could discriminate between astrocytic and neuronal NF- κ B activation. The transgenic expression of transdominant I κ B- α by a neuron-specific NSE promoter resulted in a reduction of ischemic cell death, whereas the expression of I κ B in astrocytes did not reduce the infarct size [132]. These data argue against the neuroprotective role of NF- κ B in ischemia, but it might be considered that brain reperfusion following ischemia can lead to a widespread inhibition of neuronal protein synthesis. Thus, the transcription factor NF- κ B might execute its function before brain ischemia. Here, we propose that the inhibition of neuronal NF- κ B might act in a similar way as ischemic preconditioning, thus fortifying neurons against ischemic stress.

On the other hand, under pathological conditions, the anti-apoptotic role of NF- κ B might be reversed into a pro-apoptotic function. Recent data suggest that p65 can be switched from an activator into a dominant repressor of transcription in a promoter-specific manner [133]. The p65-mediated transcription of anti-apoptotic genes through the recruitment of co-activators (histone acetyltransferases) promotes survival under physiological conditions. Under pathophysiological conditions, p65 is switched to a promoter selective dominant repressor of anti-apoptotic genes. This switch to a repressor might be mediated by the recruitment of co-repressors (histone deacetylases).

7. NF- κ B and neuronal synaptic plasticity

It has been proposed by Aryeh Routtenberg [134] that synaptic plasticity leads to information storage as the result of a synaptic dialogue. The first step would be glutamate release from the presynaptic site, followed by a modification of the presynaptic release process. This process is thought to require a retrograde messenger, which travels along the axon to switch on gene expression, in order to replenish presynaptic protein supply. We and others think that NF- κ B might be crucially involved in this important process of synaptic plasticity. In the following,

we will shortly review the evidence accumulated so far (see Fig. 2).

The capability of NF- κ B in transmitting information from active synapses to the nucleus is supported by several studies demonstrating the presence of NF- κ B in synapses [36,135,136]. Synaptosomes contain presynaptic proteins that are sealed and stabilized by the postsynaptic density. Inducible forms of NF- κ B have been found in synaptosomal preparations [36]. Low-salt extracts prepared from synaptosomes contain NF- κ B proteins, such as p50 and p65, together with I κ B- α . Synaptophysin cofractionates with NF- κ B proteins during purification. The colocalization of synaptophysin and NF- κ B proteins has also been detected in rat cerebral cortex [36]. In cortical extracts, NF- κ B DNA-binding could be activated by the detergent desoxycholate (DOC), resulting in 2 specific DNA-binding complexes with different sensitivities for DOC. Super-shifting and inhibition with recombinant I κ B- α showed a bona fide DNA-binding complex that includes the p65 and p50 subunits. Similar complexes were detected using hippocampal synaptosomal preparations [135,136]. In addition, Meberg and coworkers reported a robust increase of p65 mRNA after long-term potentiation, in vivo. It is possible that this is part of a feed-forward mechanism leading to increased DNA-binding to κ B elements during long-term potentiation. Recently, also an important influence of NF- κ B on long-term suppression of synaptic transmission was reported [137]. Purkinje cell synapses were analyzed using light microscopy and en passant synapses were found to contain NF- κ B [42]. Using electron microscopy, NF- κ B- and I κ B- α -like immunoreactivities within dendrites, including dendritic spines and postsynaptic densities, of neurons in the hippocampus and the cerebral cortex [138] were reported. With the help of an activity-specific anti-p65, antibody activated NF- κ B was detected in granule cell dendrites within the fascia dentata of rat hippocampus [139]. NF- κ B is activated in neurons by glutamate and depolarization [41,42]. In *Drosophila melanogaster*, the NF- κ B homolog Dorsal colocalizes with the I κ B homolog Cactus within the nervous system. Both proteins are detected at high levels in postsynaptic sites of glutamatergic neuromuscular junctions [140]. Thus, NF- κ B is utilized as a retrograde messenger in both pre- and postsynaptic compartments (see Fig. 2). Memory consolidation in crab also involves the activation of NF- κ B-like activity [141]. Activated NF- κ B was detected in Aplysia axons [142] and in rat [143]. In rat, traumatic brain injury first an activation of axonal NF- κ B and later activated NF- κ B is detected in neuronal nuclei. Activation could be detected as long as 1 year after brain injury [144]. As a potential injury or stress signal, sensed by synapses, in the vicinity of diffuse plaques, neurons show activated synaptic NF- κ B [145]. This might explain the activation of NF- κ B in neurons around diffuse plaques, which is lost in neurons around later plaque stages [51]. Recently, the transport of NF- κ B in living neurons was analyzed [136,146]. To enable analysis of translocation of GFP-tagged p65 in living

hippocampal neurons, a GFP tag was fused to the p65 subunit of NF- κ B and confirmed that this fusion protein (GFP-p65) retained its functionality as a transcription factor. GFP-p65 was present in the nuclei of neurons, but after overexpression, together with I κ B- α the distribution of the protein switched from nuclear to neuritic (in dendrites and axons). In glutamate-stimulated hippocampal neurons, a return of GFP-p65 from a neuritic to a nuclear distribution was observed. Meffert and colleagues reported a similar result [136]. Glutamate-induced movement of GFP-p65 was detected in hippocampal neurons. When using low amounts of GFP-p65 expression vectors, endogenous I κ B was able to keep the fusion protein in neuritic/cytoplasmic localization. Glutamate agonists could overcome the localization to neurites and activate a nuclear localization. Synaptic localization was crucially dependent on p65 [136]. In KO p65 animals, which are viable when crossed to a TNF-RI background, no synaptic NF- κ B activity was detected [136]. Interestingly, the redistribution of GFP-p65 was dependent on a functional nuclear localization sequence (NLS) [146]. This NLS-dependent retrograde transport was already described for Aplysia axons [147]. Recently, it was reported that axonal injury leads to increased retrograde transport of NLS peptides [148]. This transport was mediated by an importin α/β complex interacting with the retrograde motorprotein dynein [148]. NF- κ B activation by the neurotransmitter glutamate was identified [41,42,89,149] in cerebellar granule cells. The constitutive activity of NF- κ B was initially identified within neurons from the hippocampus and the cerebral cortex, using EMSA and immunofluorescence with an antibody specific for the activated p65 and with reporter gene assays [36,39,150–152]. It was suggested that constitutive NF- κ B activity is the result of synaptic activity [13,39,41]. The basal constitutive NF- κ B activity in neurons could be repressed by specific inhibitors, of action potential generation, glutamate receptors and L-type calcium channels [136]. The blockade of constitutive active NF- κ B was most effective through the inhibition of *N*-methyl-D-aspartate (NMDA) receptors using APV. The blockade of L-type calcium channels with nimodipine was also effective. This suggests that extracellular influx of Ca²⁺ either through NMDA receptors or via L-type Ca²⁺ channels could activate NF- κ B. Fig. 2 depicts potential pre- and postsynaptic activation mechanisms for NF- κ B. There are some evidences for the presynaptic NF- κ B activation machinery. The proteasome and ubiquitination enzymes could be detected in the presynapse [153]. Presynaptic mechanisms might involve the localized action of the IKK complex and voltage-gated Ca²⁺ channels, activated during action potential propagation. Calcium ion-mediated activation of calcium/calmodulin-dependent protein kinase II (CaMKII) is present in the presynapse [154]. Also, within the postsynapse, a potential NF- κ B activation machinery is in place. Activation by NMDA suggests a postsynaptic signaling where at least some isoforms of CamK (mainly isoform 2) have been reported to be associated with the NMDA receptor. The

autophosphorylation of α CaMKII at Thr 286 by Ca^{2+} influx through NMDA receptors [155] switches the kinase into a calcium/calmodulin-independent active status. It has been shown that CamKII can activate NF- κ B in the neurons [89,136]. The way of NF- κ B activation by CamK is not solved so far. A contribution of the proteasomal degradation machine seems likely. Indeed, an activation of the post-synaptic proteasomal degradation has been shown after the induction of neuronal activity [34]. The role of calpain remains conflicting, it seems to be active in I κ B degradation in cerebellar granule cells [156] but not in matured hippocampal cultures [136].

The activation of NF- κ B by glutamate in the cerebellum and the constitutive activity within the basal forebrain neurons could be also detected in mice containing NF- κ B reporter genes [149,150,152]. There are some reports on the failure of NF- κ B activation in neurons by glutamate [46]. A potential reason for these negative results might be the inverted U-shaped activation curve of NF- κ B (see [51] and discussion) or the sensitivity of the EMSA method which could be only used to detect activated NF- κ B in brain neurons, when isolated nuclei were prepared [39]. Another reason for these conflicting results might be the use of different culture conditions.

We conclude that NF- κ B is capable of being a signal transducer, transmitting information from, e.g., active synapses to the nucleus, in addition to its well-known role as a transcription factor. NF- κ B transduces in this manner a synaptic signal into a transcriptional event.

8. NF- κ B in learning and memory

NF- κ B might be involved in translating short-term signals from distant sites in neurites into long-term changes in gene expression, which may play a key role in plasticity, development and survival. Indeed, it has been shown that p65 $-/-$ on a TNF-RI $-/-$ background results in a severe learning deficit [136]. There are two other mouse models where NF- κ B was repressed via tetracycline-regulated expression of transdominant negative I κ B: (1) a model with CamKII promoter-driven expression of tTA in basal forebrain neurons [151]; (2) a model with prion promoter-driven tTA expression in neurons and glia by WC Greene's group (see [157]). Both have a modulation of learning and memory. For an extensive discussion, see the recent comprehensive review by [124]. In accordance with Meffert and coworkers, we have found that the repression of NF- κ B by I κ B in neurons resulted in behavioral deficits and a reduction in LTP and LTD induction. These effects could be correlated with a strongly reduced CREB phosphorylation [172]. On the other hand, I κ B expression driven by the prion promoter-expressed tTA resulted in enhanced learning in older animals (WC Greene and coworkers, see [124]). These might be two sides of the same coin. On one side, a repression of neuroinflammation

in elder age via the inhibition of pathological NF- κ B hyperactivation might enhance learning. On the other side, neuronal NF- κ B at physiological levels is needed for learning.

9. NF- κ B in neuroprotection and neurodegeneration

Initial reports of the neuroprotective role of TNF [11] were followed by a neuroprotective role for NF- κ B. DNA decoy with NF- κ B binding sites did compete in cultured neurons with NF- κ B activity induced by TNF and did abolish neuroprotection [54,158]. Similarly, survival of adult sensory neurons is dependent on TNF-mediated NF- κ B activation [159]. In cerebellar granule cells, which have a low basal NF- κ B activity [41], an inverted U-shaped dose response for TNF-mediated NF- κ B activation could be detected. Also an inverted U-shaped protection curve was also reported, when TNF-mediated protection against NMDA excitotoxicity [12] was analyzed. Initially it was shown that TNF-mediated signaling via NF- κ B could protect neurons against excitotoxic stress and against neurotoxic amyloid β [11,51,54]. The main component of amyloid β consists of a 40- to 43- amino acid peptide, A β [160], which is derived from a proteolytic cleavage of a family of ubiquitously expressed trans-membrane proteins,




Fig. 3. Model for NF- κ B homeostasis. Ongoing synaptic transmission might provide the permanent activation signal for constitutively active NF- κ B in neurons. A physiological activation state could be maintained via an autoregulatory loop to the expression of NF- κ B inhibiting proteins such as I κ B family members or de-ubiquitinating enzymes, e.g., CYLD or A20. A perturbation of this physiological NF- κ B activation might be pathological with the resulting cell death. The p65-mediated transcription of anti-apoptotic genes through the recruitment of co-activators (histone acetyltransferases) promotes survival under physiological conditions. Under pathophysiological conditions, p65 could be switched to a promoter selective dominant repressor of anti-apoptotic genes. This switch to a repressor might be mediated by the recruitment of co-repressors (histone deacetylases). This might be an additional mechanism for generating pathological low NF- κ B level.

named amyloid precursor proteins (APP) [161]. Under normal conditions, the most abundant A β isoform is A β 1–40; however, much of the fibrillar A β is composed of the longer A β 1–42 isoform [162]. Initially A β 1–42 is deposited in the form of an immature diffuse plaque, with no or little neuritic dystrophy [162]. In control subjects, NF- κ B activation could be detected around all plaque types including diffuse and neuritic plaque stages (primitive, classical and compact) [51]. In Alzheimer patients, NF- κ B activation around all analyzed plaque stages was reduced. Plaques were classified as diffuse and neuritic (primitive, classical and compact); see Kaltschmidt et al. [51].

In respect of this, it is noteworthy that the NF- κ B signaling pathway is one of the major neuroprotective pathways identified to be protective against Alzheimer's disease (see [163] for a detailed discussion). A reduction of this protective NF- κ B activation within Alzheimer patients' brains around late plaque stages might be one of the reasons for increased neurodegeneration [51].

A low dose of A β was able to activate NF- κ B and to protect against a high cytotoxic dose of A β . This led to the discovery of an essential role for NF- κ B in preconditioning [55,164,165]. Preconditioning describes an old observation worded by Paracelsus as "Alle Ding' sind Gift und nichts ohn' Gift; allein die Dosis macht, das ein Ding' kein Gift ist" (all things are poisons, only the dose makes the poison). In this line, a toxin in a low dose could activate a cellular response program, which later on protects against a high dose of toxin. We think that the mechanisms might be similar to a process described by David Baltimore as intracellular immunization [166] against virus infection. The preconditioning effect of NF- κ B could be completely abolished by the overexpression of transdominant I κ B- α . On the other hand, there are toxic stimuli such as staurosporine where repression of NF- κ B activation was protective [115]. This might be a general concept, which we first described in non-neuronal cells: The nature of the apoptotic stimulus dictates the pro- or anti-apoptotic action of NF- κ B [167]. Genetic evidence suggests that constitutive NF- κ B activity is essential for neuronal survival [152]. Similarly, pharmacological repression of the IKK complex ([168]; B. Kaltschmidt et al., unpublished) results in neuronal death. The transgenic overexpression of transdominant I κ B- α sensitizes neurons against excitotoxic lesions [151]. NF- κ B activation protects neurons against A β toxicity [51,54]. This might be a neuroprotective mechanism which is perturbed during Alzheimer disease [163]. On the other hand, several diseases where NF- κ B hyperactivation is disease promoting are described. Examples include ischemia [82], Parkinson [169] and conditions where NF- κ B-dependent p53 transcription mediates neuronal death [168]. The p53-mediated cell death seems to rely on a repression of NF- κ B activity [170] in some paradigms.

To sort this dilemma out, an optimal activation hypothesis (Fig. 3) is proposed similar to the NMDA receptor activation [171]. Too low activation of NF- κ B in neurons is

disastrous, as is too high activation (hyperactivation). The optimal basal constitutive level in neurons is maintained by synaptic activity, which activates NF- κ B and is repressed by one of its target genes I κ B or CYLD in an auto-regulatory fashion (see Fig. 3). A potential protective role of NF- κ B in neurons should be discriminated from a potential degenerative role in glia.

Acknowledgements

The work in our laboratory was supported, in part, by grants from the Deutsche Forschungsgemeinschaft (DFG), the Volkswagen-Stiftung and the Land NRW.

References

- [1] G. Nabel, D. Baltimore, An inducible transcription factor activates expression of human immunodeficiency virus in T cells, *Nature* 326 (1987) 711–713.
- [2] M.L. Schmitz, I. Mattioli, H. Buss, M. Kracht, NF-kappaB: a multifaceted transcription factor regulated at several levels, *Chem-BioChem* 5 (2004) 1348–1358.
- [3] G. Bonizzi, M. Karin, The two NF-kappaB activation pathways and their role in innate and adaptive immunity, *Trends Immunol.* 25 (2004) 280–288.
- [4] A.J. Bruce, W. Boling, M.S. Kindy, J. Peschon, P.J. Kraemer, M.K. Carpenter, F.W. Holtsberg, M.P. Mattson, Altered neuronal and microglial responses to excitotoxic and ischemic brain injury in mice lacking TNF receptors, *Nat. Med.* 2 (1996) 788–794.
- [5] M.A. Collart, P. Baeuerle, P. Vassalli, Regulation of tumor necrosis factor alpha transcription in macrophages: involvement of four kappa B-like motifs and of constitutive and inducible forms of NF-kappa B, *Mol. Cell. Biol.* 10 (1990) 1498–1506.
- [6] R. Kolesnick, D.W. Golde, The sphingomyelin pathway in tumor necrosis factor and interleukin-1 signaling, *Cell* 77 (1994) 325–328.
- [7] F. Henkler, H. Wajant, in: M.L. Schmitz, S. Bacher (Eds.), *Mechanisms of Signal Transduction and Inducible Gene Expression*, Research Signpost, Kerala, India, 2004, pp. 97–119.
- [8] L. Yang, K. Lindholm, Y. Konishi, R. Li, Y. Shen, Target depletion of distinct tumor necrosis factor receptor subtypes reveals hippocampal neuron death and survival through different signal transduction pathways, *J. Neurosci.* 22 (2002) 3025–3032.
- [9] L. Marchetti, M. Klein, K. Schlett, K. Pfizenmaier, U.L. Eisel, Tumor necrosis factor (TNF)-mediated neuroprotection against glutamate-induced excitotoxicity is enhanced by *N*-methyl-D-aspartate receptor activation. Essential role of a TNF receptor 2-mediated phosphatidylinositol 3-kinase-dependent NF-kappa B pathway, *J. Biol. Chem.* 279 (2004) 32869–32881.
- [10] O. Micheau, J. Tschopp, Induction of TNF receptor I-mediated apoptosis via two sequential signaling complexes, *Cell* 114 (2003) 181–190.
- [11] B. Cheng, S. Christakos, M.P. Mattson, Tumor necrosis factors protect neurons against metabolic–excitotoxic insults and promote maintenance of calcium homeostasis, *Neuron* 12 (1994) 139–153.
- [12] N.G. Carlson, W.A. Wieggl, J. Chen, A. Bacchi, S.W. Rogers, L.C. Gahring, Inflammatory cytokines IL-1 alpha, IL-1 beta, IL-6, and TNF-alpha impart neuroprotection to an excitotoxin through distinct pathways, *J. Immunol.* 163 (1999) 3963–3968.
- [13] L.A.J. O'Neill, C. Kaltschmidt, NF- κ B: a crucial transcription factor for glial and neuronal cell function, *Trends Neurosci.* 20 (1997) 252–258.

- [14] J.L. Ducut Sigala, V. Bottero, D.B. Young, A. Shevchenko, F. Mercurio, I.M. Verma, Activation of transcription factor NF-kappaB requires ELKS, an IkappaB kinase regulatory subunit, *Science* 304 (2004) 1963–1967.
- [15] M. Delhase, M. Hayakawa, Y. Chen, M. Karin, Positive and negative regulation of IkappaB kinase activity through IKKbeta subunit phosphorylation, *Science* 284 (1999) 309–313.
- [16] J. Yang, Y. Lin, Z. Guo, J. Cheng, J. Huang, L. Deng, W. Liao, Z. Chen, Z. Liu, B. Su, The essential role of MEKK3 in TNF-induced NF-kappaB activation, *Nat. Immunol.* 2 (2001) 620–624.
- [17] S. Tegethoff, J. Behlke, C. Scheidereit, Tetrameric oligomerization of IkappaB kinase gamma (IKKgamma) is obligatory for IKK complex activity and NF-kappaB activation, *Mol. Cell. Biol.* 23 (2003) 2029–2041.
- [18] F. Agou, G. Courtois, J. Chiaravalli, F. Baleux, Y.M. Coic, F. Traincard, A. Israel, M. Veron, Inhibition of NF-kappa B activation by peptides targeting NF-kappa B essential modulator (nemo) oligomerization, *J. Biol. Chem.* 279 (2004) 54248–54257.
- [19] R. Weil, K. Schwamborn, A. Alcover, C. Bessia, V. Di Bartolo, A. Israel, Induction of the NF-kappaB cascade by recruitment of the scaffold molecule NEMO to the T cell receptor, *Immunity* 18 (2003) 13–26.
- [20] G. Chen, P. Cao, D.V. Goeddel, TNF-induced recruitment and activation of the IKK complex require Cdc37 and Hsp90, *Mol. Cell* 9 (2002) 401–410.
- [21] A. Kovalenko, C. Chable-Bessia, G. Cantarella, A. Israel, D. Wallach, G. Courtois, The tumour suppressor CYLD negatively regulates NF-kappaB signalling by deubiquitination, *Nature* 424 (2003) 801–805.
- [22] E. Trompouki, E. Hatzivassiliou, T. Tschritzis, H. Farmer, A. Ashworth, G. Mosialos, CYLD is a deubiquitinating enzyme that negatively regulates NF-kappaB activation by TNFR family members, *Nature* 424 (2003) 793–796.
- [23] K. Heyninck, R. Beyaert, A20 inhibits NF-kappaB activation by dual ubiquitin-editing functions, *Trends Biochem. Sci.* 30 (2005) 1–4.
- [24] A. Ciechanover, A.L. Schwartz, The ubiquitin system: pathogenesis of human diseases and drug targeting, *Biochim. Biophys. Acta* 1695 (2004) 3–17.
- [25] A. Yaron, A. Hatzubai, M. Davis, I. Lavon, S. Amit, A.M. Manning, J.S. Andersen, M. Mann, F. Mercurio, Y. Ben-Neriah, Identification of the receptor component of the IkappaBalpha-ubiquitin ligase, *Nature* 396 (1998) 590–594.
- [26] M. Hayakawa, H. Miyashita, I. Sakamoto, M. Kitagawa, H. Tanaka, H. Yasuda, M. Karin, K. Kikugawa, Evidence that reactive oxygen species do not mediate NF-kappaB activation, *EMBO J.* 22 (2003) 3356–3366.
- [27] M.D. Jacobs, S.C. Harrison, Structure of an IkappaBalpha/NF-kappaB complex, *Cell* 95 (1998) 749–758.
- [28] T. Huxford, D.B. Huang, S. Malek, G. Ghosh, The crystal structure of the IkappaBalpha/NF-kappaB complex reveals mechanisms of NF-kappaB inactivation, *Cell* 95 (1998) 759–770.
- [29] S. Malek, T. Huxford, G. Ghosh, Ikappa Balpha functions through direct contacts with the nuclear localization signals and the DNA binding sequences of NF-kappaB, *J. Biol. Chem.* 273 (1998) 25427–25435.
- [30] R. Fagerlund, L. Kinnunen, M. Kohler, I. Julkunen, K. Melen, NF-kappa B is transported into the nucleus by importin alpha 3 and importin alpha 4, *J. Biol. Chem.* 280 (16) (2005) 15942–15951.
- [31] S. Sacconi, I. Marazzi, A.A. Beg, G. Natoli, Degradation of promoter-bound p65/RelA is essential for the prompt termination of the nuclear factor kappaB response, *J. Exp. Med.* 200 (2004) 107–113.
- [32] M. Rape, S. Jentsch, Productive RUPTure: activation of transcription factors by proteasomal processing, *Biochim. Biophys. Acta* 1695 (2004) 209–213.
- [33] L. Lin, G.N. DeMartino, W.C. Greene, Cotranslational biogenesis of NF-kappaB p50 by the 26S proteasome, *Cell* 92 (1998) 819–828.
- [34] M.D. Ehlers, Activity level controls postsynaptic composition and signaling via the ubiquitin-proteasome system, *Nat. Neurosci.* 6 (2003) 231–242.
- [35] S.T. Whiteside, A. Israel, I kappa B proteins: structure, function and regulation, *Semin. Cancer Biol.* 8 (1997) 75–82.
- [36] C. Kaltschmidt, B. Kaltschmidt, P.A. Baeuerle, Brain synapses contain inducible forms of the transcription factor NF-kB, *Mech. Dev.* 43 (1993) 135–147.
- [37] G. Bakalkin, T. Yakovleva, L. Terenius, NF-kappa B-like factors in the murine brain. Developmentally-regulated and tissue-specific expression, *Brain Res. Mol. Brain Res.* 20 (1993) 137–146.
- [38] A. Rattner, M. Korner, M.D. Walker, Y. Citri, NF-kappa B activates the HIV promoter in neurons, *EMBO J.* 12 (1993) 4261–4267.
- [39] C. Kaltschmidt, B. Kaltschmidt, H. Neumann, H. Wekerle, P.A. Baeuerle, Constitutive NF-kB activity in neurons, *Mol. Cell. Biol.* 14 (1994) 3981–3992.
- [40] C. Kaltschmidt, B. Kaltschmidt, T. Henkel, H. Stockinger, P.A. Baeuerle, Selective recognition of the activated form of transcription factor NF-kappa B by a monoclonal antibody, *Biol. Chem. Hoppe Seyler* 376 (1995) 9–16.
- [41] C. Kaltschmidt, B. Kaltschmidt, P.A. Baeuerle, Stimulation of ionotropic glutamate receptors activates transcription factor NF-kB in primary neurons, *Proc. Natl. Acad. Sci. U. S. A.* 92 (1995) 9618–9622.
- [42] L. Guerrini, F. Blasi, D.S. Denis, Synaptic activation of NF-kB by glutamate in cerebellar granule neurons in vitro, *Proc. Natl. Acad. Sci. U. S. A.* 92 (1995) 9077–9081.
- [43] B. Kaltschmidt, M. Uhrek, B. Volk, P.A. Baeuerle, C. Kaltschmidt, Transcription factor NF-kB is activated in primary neurons by amyloid β peptides and in neurons surrounding early plaques from patients with Alzheimer disease, *Proc. Natl. Acad. Sci. U. S. A.* 94 (1997) 2642–2647.
- [44] M. Koerner, A. Rattner, F. Mauxion, R. Sen, Y. Citri, A brain-specific transcription activator, *Neuron* 3 (1989) 563–572.
- [45] K. Cauley, I.M. Verma, Kappa B enhancer-binding complexes that do not contain NF-kappa B are developmentally regulated in mammalian brain, *Proc. Natl. Acad. Sci. U. S. A.* 91 (1994) 390–394.
- [46] A.M. Moerman, X. Mao, M.M. Lucas, S.W. Barger, Characterization of a neuronal kappaB-binding factor distinct from NF-kappaB, *Brain Res. Mol. Brain Res.* 67 (1999) 303–315.
- [47] E. Bourke, E.J. Kennedy, P.N. Moynagh, Loss of Ikappa B-beta is associated with prolonged NF-kappa B activity in human glial cells, *J. Biol. Chem.* 275 (2000) 39996–40002.
- [48] I. Kemler, A. Fontana, Role of IkappaBalpha and IkappaB-beta in the biphasic nuclear translocation of NF-kappaB in TNFalpha-stimulated astrocytes and in neuroblastoma cells, *Glia* 26 (1999) 212–220.
- [49] B.D. Carter, C. Kaltschmidt, B. Kaltschmidt, N. Offenhauser, R. Bohm-Matthaei, P.A. Baeuerle, Y.A. Barde, Selective activation of NF-kB by nerve growth factor through the neurotrophin receptor p75, *Science* 272 (1996) 542–545.
- [50] K. Nakajima, Y. Kikuchi, E. Ikoma, S. Honda, M. Ishikawa, Y. Liu, S. Kohsaka, Neurotrophins regulate the function of cultured microglia, *Glia* 24 (1998) 272–289.
- [51] B. Kaltschmidt, M. Uhrek, H. Wellmann, B. Volk, C. Kaltschmidt, Inhibition of NF-kB potentiates amyloid- β -mediated neuronal apoptosis, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 9409–9414.
- [52] R.H. Lipsky, K. Xu, D. Zhu, C. Kelly, A. Terhakopian, A. Novelli, A.M. Marini, Nuclear factor kappaB is a critical determinant in N-methyl-D-aspartate receptor-mediated neuroprotection, *J. Neurochem.* 78 (2001) 254–264.
- [53] P.D. Drew, M. Loneran, M.E. Goldstein, L.A. Lampson, K. Ozato, D.E. McFarlin, Regulation of MHC class I and beta 2-microglobulin gene expression in human neuronal cells. Factor binding to conserved cis-acting regulatory sequences correlates with expression of the genes, *J. Immunol.* 150 (1993) 3300–3310.
- [54] S.W. Barger, D. Horster, K. Furukawa, Y. Goodman, J. Kriegelstein, M.P. Mattson, Tumor necrosis factors alpha and beta protect neurons

- against amyloid beta-peptide toxicity: evidence for involvement of a kappa B-binding factor and attenuation of peroxide and Ca^{2+} accumulation, *Proc. Natl. Acad. Sci. U. S. A.* 92 (1995) 9328–9332.
- [55] B. Kaltschmidt, M. Uhrek, H. Wellmann, B. Volk, C. Kaltschmidt, Inhibition of NF-kappaB potentiates amyloid beta-mediated neuronal apoptosis, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 9409–9414.
- [56] S.M. Sparacio, Y. Zhang, J. Vilcek, E.N. Benveniste, Cytokine regulation of interleukin-6 gene expression in astrocytes involves activation of an NF-kappa B-like nuclear protein, *J. Neuroimmunol.* 39 (1992) 231–242.
- [57] S.J. Lee, T. Zhou, C. Choi, Z. Wang, E.N. Benveniste, Differential regulation and function of Fas expression on glial cells, *J. Immunol.* 164 (2000) 1277–1285.
- [58] M. Grilli, F. Goffi, M. Memo, P. Spano, Interleukin-1beta and glutamate activate the NF-kappaB/Rel binding site from the regulatory region of the amyloid precursor protein gene in primary neuronal cultures, *J. Biol. Chem.* 271 (1996) 15002–15007.
- [59] S.W. Barger, M.P. Mattson, Induction of neuroprotective kappa B-dependent transcription by secreted forms of the Alzheimer's beta-amyloid precursor, *Mol. Brain Res.* 40 (1996) 116–126.
- [60] S.W. Barger, A.D. Harmon, Microglial activation by Alzheimer amyloid precursor protein and modulation by apolipoprotein E, *Nature* 388 (1997) 878–881.
- [61] C. Behl, J.B. Davis, R. Lesley, D. Schubert, Hydrogen peroxide mediates amyloid beta protein toxicity, *Cell* 77 (1994) 817–827.
- [62] C. Bonaiuto, P.P. McDonald, F. Rossi, M.A. Cassatella, Activation of nuclear factor-kappa B by beta-amyloid peptides and interferon-gamma in murine microglia, *J. Neuroimmunol.* 77 (1997) 51–56.
- [63] D. Ferrari, S. Wesselborg, M. Bauer, O.K. Schulze, Extracellular ATP activates transcription factor NF-kappaB through the P2Z purinoreceptor by selectively targeting NF-kappaB p65, *J. Cell Biol.* 139 (1997) 1635–1643.
- [64] G.R. John, J.E. Simpson, M.N. Woodroffe, S.C. Lee, C.F. Brosnan, Extracellular nucleotides differentially regulate interleukin-1beta signaling in primary human astrocytes: implications for inflammatory gene expression, *J. Neurosci.* 21 (2001) 4134–4142.
- [65] R. Basheer, D.G. Rainnie, T. Porkka-Heiskanen, V. Ramesh, R.W. McCarley, Adenosine, prolonged wakefulness, and A1-activated NF-kappaB DNA binding in the basal forebrain of the rat, *Neuroscience* 104 (2001) 731–739.
- [66] M.K. Bauer, K. Lieb, K. Schulze-Osthoff, M. Berger, P.J. Gebicke-Haerter, J. Bauer, B.L. Fiebich, Expression and regulation of cyclooxygenase-2 in rat microglia, *Eur. J. Biochem.* 243 (1997) 726–731.
- [67] G. Pistrutto, O. Franzese, G. Pozzoli, C. Mancuso, G. Tringali, P. Preziosi, P. Navarra, Bacterial lipopolysaccharide increases prostaglandin production by rat astrocytes via inducible cyclo-oxygenase: evidence for the involvement of nuclear factor kappaB, *Biochem. Biophys. Res. Commun.* 263 (1999) 570–574.
- [68] M. Digiacylioglu, S.A. Lipton, Erythropoietin-mediated neuroprotection involves cross-talk between Jak2 and NF-kappaB signalling cascades, *Nature* 412 (2001) 641–647.
- [69] M. Schwaninger, S. Sallmann, N. Petersen, A. Schneider, S. Prinz, T.A. Libermann, M. Spranger, Bradykinin induces interleukin-6 expression in astrocytes through activation of nuclear factor-kappaB, *J. Neurochem.* 73 (1999) 1461–1466.
- [70] Y. Han, T. He, D.R. Huang, C.A. Pardo, R.M. Ransohoff, TNF-alpha mediates SDF-1 alpha-induced NF-kappa B activation and cytotoxic effects in primary astrocytes, *J. Clin. Invest.* 108 (2001) 425–435.
- [71] O. Zelenai, B.D. Schlag, G.E. Gochenauer, R. Ganel, W. Song, J.S. Beesley, J.B. Grinspan, J.D. Rothstein, M.B. Robinson, Epidermal growth factor receptor agonists increase expression of glutamate transporter GLT-1 in astrocytes through pathways dependent on phosphatidylinositol 3-kinase and transcription factor NF-kappaB, *Mol. Pharmacol.* 57 (2000) 667–678.
- [72] K.L. Jin, X.O. Mao, D.A. Greenberg, Vascular endothelial growth factor rescues HN33 neural cells from death induced by serum withdrawal, *J. Mol. Neurosci.* 14 (2000) 197–203.
- [73] J.N. Wood, Regulation of NF-kappa B activity in rat dorsal root ganglia and PC12 cells by tumour necrosis factor and nerve growth factor, *Neurosci. Lett.* 192 (1995) 41–44.
- [74] S.O. Yoon, P. Casaccia-Bonnel, B. Carter, M.V. Chao, Competitive signaling between TrkA and p75 nerve growth factor receptors determines cell survival, *J. Neurosci.* 18 (1998) 3273–3281.
- [75] G. Middleton, M. Hamanoue, Y. Enokido, S. Wyatt, D. Pennica, E. Jaffray, R.T. Hay, A.M. Davies, Cytokine-induced nuclear factor kappa B activation promotes the survival of developing neurons, *J. Cell Biol.* 148 (2000) 325–332.
- [76] S. Heck, F. Lezoualc'h, S. Engert, C. Behl, Insulin-like growth factor-1-mediated neuroprotection against oxidative stress is associated with activation of nuclear factor kappaB, *J. Biol. Chem.* 274 (1999) 9828–9835.
- [77] G.W. Glazner, S. Camandola, M.P. Mattson, Nuclear factor-kappaB mediates the cell survival-promoting action of activity-dependent neurotrophic factor peptide-9, *J. Neurochem.* 75 (2000) 101–108.
- [78] T. Yabe, D. Wilson, J.P. Schwartz, NF-kappaB activation is required for the neuroprotective effects of pigment epithelium-derived factor (PEDF) on cerebellar granule neurons, *J. Biol. Chem.* 276 (2001) 43313–43319.
- [79] U. Vollgraf, M. Wegner, C. Richter-Landsberg, Activation of AP-1 and nuclear factor-kappaB transcription factors is involved in hydrogen peroxide-induced apoptotic cell death of oligodendrocytes, *J. Neurochem.* 73 (1999) 2501–2509.
- [80] H. Pyo, E. Joe, S. Jung, S.H. Lee, I. Jou, Gangliosides activate cultured rat brain microglia, *J. Biol. Chem.* 274 (1999) 34584–34589.
- [81] D. Daily, A. Vlamis-Gardikas, D. Offen, L. Mittelman, E. Melamed, A. Holmgren, A. Barzilay, Glutaredoxin protects cerebellar granule neurons from dopamine-induced apoptosis by dual activation of the ras-phosphoinositide 3-kinase and jun N-terminal kinase pathways, *J. Biol. Chem.* 276 (2001) 21618–21626.
- [82] A. Schneider, A. Martin-Villalba, F. Weih, J. Vogel, T. Wirth, M. Schwaninger, NF-kappaB is activated and promotes cell death in focal cerebral ischemia, *Nat. Med.* 5 (1999) 554–559.
- [83] A. Bierhaus, K.M. Haslbeck, P.M. Humpert, B. Liliensiek, T. Dehmer, M. Morcos, A.A. Sayed, M. Andrassy, S. Schiekofe, J.G. Schneider, J.B. Schulz, D. Heuss, B. Neundorfer, S. Dierl, J. Huber, H. Tritschler, A.M. Schmidt, M. Schwaninger, H.U. Haering, E. Schleicher, M. Kasper, D.M. Stern, B. Arnold, P.P. Nawroth, Loss of pain perception in diabetes is dependent on a receptor of the immunoglobulin superfamily, *J. Clin. Invest.* 114 (2004) 1741–1751.
- [84] S. Tanaka, M. Takehashi, N. Matoh, S. Iida, T. Suzuki, S. Futaki, H. Hamada, E. Maslia, Y. Sugiura, K. Ueda, Generation of reactive oxygen species and activation of NF-kappaB by non-Abeta component of Alzheimer's disease amyloid, *J. Neurochem.* 82 (2002) 305–315.
- [85] P. Bartholoma, N. Erlandsson, K. Kaufmann, O.G. Rossler, B. Baumann, T. Wirth, K.M. Giehl, G. Thiel, Neuronal cell death induced by antidepressants: lack of correlation with Egr-1, NF-kappa B and extracellular signal-regulated protein kinase activation, *Biochem. Pharmacol.* 63 (2002) 1507–1516.
- [86] M. Pizzi, F. Goffi, F. Boroni, M. Benarese, S.E. Perkins, H.C. Liou, P. Spano, Opposing roles for NF-kappa B/Rel factors p65 and c-Rel in the modulation of neuron survival elicited by glutamate and interleukin-1beta, *J. Biol. Chem.* 277 (2002) 20717–20723.
- [87] D.R. Brown, R.S. Nicholas, L. Canevari, Lack of prion protein expression results in a neuronal phenotype sensitive to stress, *J. Neurosci. Res.* 67 (2002) 211–224.
- [88] N.E. Savaskan, A.U. Brauer, M. Kuhbacher, I.Y. Eyupoglu, A. Kyriakopoulos, O. Ninnemann, D. Behne, R. Nitsch, Selenium deficiency increases susceptibility to glutamate-induced excitotoxicity, *FASEB J.* 17 (2003) 112–114.

- [89] A. Lilienbaum, A. Israel, From calcium to NF-kappa B signaling pathways in neurons, *Mol. Cell. Biol.* 23 (2003) 2680–2698.
- [90] G.A. de Erausquin, K. Hyrc, D.A. Dorsey, D. Mamah, M. Dokucu, D.H. Masco, T. Walton, K. Dikranian, M. Soriano, J.M. Garcia Verdugo, M.P. Goldberg, L.L. Dugan, Nuclear translocation of nuclear transcription factor-kappa B by alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors leads to transcription of p53 and cell death in dopaminergic neurons, *Mol. Pharmacol.* 63 (2003) 784–790.
- [91] M.A. Burke, M. Bothwell, p75 neurotrophin receptor mediates neurotrophin activation of NF-kappa B and induction of iNOS expression in P19 neurons, *J. Neurobiol.* 55 (2003) 191–203.
- [92] J.C. Nickols, W. Valentine, S. Kanwal, B.D. Carter, Activation of the transcription factor NF-kappaB in Schwann cells is required for peripheral myelin formation, *Nat. Neurosci.* 6 (2003) 161–167.
- [93] I. Charalampopoulos, C. Tsatsanis, E. Dermitzaki, V.I. Alexaki, E. Castanas, A.N. Margioris, A. Gravanis, Dehydroepiandrosterone and allopregnanolone protect sympathoadrenal medulla cells against apoptosis via antiapoptotic Bcl-2 proteins, *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 8209–8214.
- [94] J.A. Brandt, L. Churchill, A. Rehman, G. Ellis, S. Memet, A. Israel, J.M. Krueger, Sleep deprivation increases the activation of nuclear factor kappa B in lateral hypothalamic cells, *Brain Res.* 1004 (2004) 91–97.
- [95] C. Sole, X. Dolcet, M.F. Segura, H. Gutierrez, M.T. Diaz-Meco, R. Gozzelino, D. Sanchis, J.R. Bayascas, C. Gallego, J. Moscat, A.M. Davies, J.X. Comella, The death receptor antagonist FAIM promotes neurite outgrowth by a mechanism that depends on ERK and NF-kappa B signaling, *J. Cell Biol.* 167 (2004) 479–492.
- [96] Y. Zhu, C. Culmsee, S. Klumpp, J. Kriegstein, Neuroprotection by transforming growth factor-beta1 involves activation of nuclear factor-kappaB through phosphatidylinositol-3-OH kinase/Akt and mitogen-activated protein kinase-extracellular-signal regulated kinase 1,2 signaling pathways, *Neuroscience* 123 (2004) 897–906.
- [97] M. Arsur, M. Wu, G.E. Sonenshein, TGF beta 1 inhibits NF-kappa B/Rel activity inducing apoptosis of B cells: transcriptional activation of I kappa B alpha, *Immunity* 5 (1996) 31–40.
- [98] S. Camandola, G. Poli, M.P. Mattson, The lipid peroxidation product 4-hydroxy-2,3-nonenal inhibits constitutive and inducible activity of nuclear factor kappa B in neurons, *Brain Res. Mol. Brain Res.* 85 (2000) 53–60.
- [99] F. Pousset, R. Dantzer, K.W. Kelley, P. Parnet, Interleukin-1 signaling in mouse astrocytes involves Akt: a study with interleukin-4 and IL-10, *Eur. Cytokine Netw.* 11 (2000) 427–434.
- [100] A. Bachis, A.M. Colangelo, S. Vicini, P.P. Doe, M.A. De Bernardi, G. Brooker, I. Mocchetti, Interleukin-10 prevents glutamate-mediated cerebellar granule cell death by blocking caspase-3-like activity, *J. Neurosci.* 21 (2001) 3104–3112.
- [101] L.C. Ehrlich, S. Hu, P.K. Peterson, C.C. Chao, IL-10 down-regulates human microglial IL-8 by inhibition of NF-kappaB activation, *NeuroReport* 9 (1998) 1723–1726.
- [102] S. Hu, C.C. Chao, L.C. Ehrlich, W.S. Sheng, R.L. Sutton, G.L. Rockswold, P.K. Peterson, Inhibition of microglial cell RANTES production by IL-10 and TGF-beta, *J. Leukoc. Biol.* 65 (1999) 815–821.
- [103] B. Kaltschmidt, C. Kaltschmidt, DNA array analysis of the developing rat cerebellum: transforming growth factor-beta2 inhibits constitutively activated NF-kappaB in granule neurons, *Mech. Dev.* 101 (2001) 11–19.
- [104] I. Ginis, J.M. Hallenbeck, J. Liu, M. Spatz, R. Jaiswal, E. Shohami, Tumor necrosis factor and reactive oxygen species cooperative cytotoxicity is mediated via inhibition of NF-kappaB, *Mol. Med.* 6 (2000) 1028–1041.
- [105] H. Togashi, M. Sasaki, E. Frohman, E. Taira, R.R. Ratan, T.M. Dawson, V.L. Dawson, Neuronal (type I) nitric oxide synthase regulates nuclear factor kappaB activity and immunologic (type II) nitric oxide synthase expression, *Proc. Natl. Acad. Sci. U. S. A.* 94 (1997) 2676–2680.
- [106] M. Colasanti, T. Persichini, Nitric oxide: an inhibitor of NF-kappaB/Rel system in glial cells, *Brain Res. Bull.* 52 (2000) 155–161.
- [107] S. Braun, W. Liebetrau, B. Berning, C. Behl, Dexamethasone-enhanced sensitivity of mouse hippocampal HT22 cells for oxidative stress is associated with the suppression of nuclear factor-kappaB, *Neurosci. Lett.* 295 (2000) 101–104.
- [108] F. Lezoualc'h, M. Sparapani, C. Behl, N-acetyl-serotonin (normelatonin) and melatonin protect neurons against oxidative challenges and suppress the activity of the transcription factor NF-kappaB, *J. Pineal Res.* 24 (1998) 168–178.
- [109] F. Lezoualc'h, S. Engert, B. Berning, C. Behl, Corticotropin-releasing hormone-mediated neuroprotection against oxidative stress is associated with the increased release of non-amyloidogenic amyloid beta precursor protein and with the suppression of nuclear factor-kappaB, *Mol. Endocrinol.* 14 (2000) 147–159.
- [110] L. Acarin, B. Gonzalez, B. Castellano, Stat3 and NFkappaB glial expression after excitotoxic damage to the postnatal brain, *NeuroReport* 9 (1998) 2869–2873.
- [111] D. Stephenson, T. Yin, E.B. Smalstig, M.A. Hsu, J. Panetta, S. Little, J. Clemens, Transcription factor nuclear factor-kappa B is activated in neurons after focal cerebral ischemia, *J. Cereb. Blood Flow Metab.* 20 (2000) 592–603.
- [112] P.T. Massa, Specific suppression of major histocompatibility complex class I and class II genes in astrocytes by brain-enriched gangliosides, *J. Exp. Med.* 178 (1993) 1357–1363.
- [113] C. Behl, Vitamin E protects neurons against oxidative cell death in vitro more effectively than 17-beta estradiol and induces the activity of the transcription factor NF-kappaB, *J. Neural Transm.* 107 (2000) 393–407.
- [114] E. Juttler, I. Potrovita, V. Tarabin, S. Prinz, T. Dong-Si, G. Fink, M. Schwaninger, The cannabinoid dexamabinol is an inhibitor of the nuclear factor-kappa B (NF-kappa B), *Neuropharmacology* 47 (2004) 580–592.
- [115] B. Kaltschmidt, M. Heinrich, C. Kaltschmidt, Stimulus-dependent activation of NF-kappaB specifies apoptosis or neuroprotection in cerebellar granule cells, *Neuromol. Med.* 2 (2002) 299–309.
- [116] S.K. Sharma, E.C. Carlson, M. Ebadi, Neuroprotective actions of Selegiline in inhibiting 1-methyl, 4-phenyl, pyridinium ion (MPP+)-induced apoptosis in SK-N-SH neurons, *J. Neurocytol.* 32 (2003) 329–343.
- [117] E.M. Halvorsen, J. Dennis, P. Keeney, T.W. Sturgill, J.B. Tuttle, J.B. Bennett Jr., Methylpyridinium (MPP+)- and nerve growth factor-induced changes in pro- and anti-apoptotic signaling pathways in SH-SY5Y neuroblastoma cells, *Brain Res.* 952 (2002) 98–110.
- [118] M.J. Wang, W.W. Lin, H.L. Chen, Y.H. Chang, H.C. Ou, J.S. Kuo, J.S. Hong, K.C. Jeng, Silymarin protects dopaminergic neurons against lipopolysaccharide-induced neurotoxicity by inhibiting microglia activation, *Eur. J. Neurosci.* 16 (2002) 2103–2112.
- [119] C.A. Kassed, T.L. Butler, M.T. Navidomskis, M.N. Gordon, D. Morgan, K.R. Pennypacker, Mice expressing human mutant presenilin-1 exhibit decreased activation of NF-kappaB p50 in hippocampal neurons after injury, *Brain Res. Mol. Brain Res.* 110 (2003) 152–157.
- [120] R. Amodio, C. De Ruvo, A. Sacchetti, A. Di Santo, N. Martelli, V. Di Matteo, R. Lorenzet, A. Poggi, D. Rotilio, M. Cacchio, E. Esposito, Caffeic acid phenethyl ester blocks apoptosis induced by low potassium in cerebellar granule cells, *Int. J. Dev. Neurosci.* 21 (2003) 379–389.
- [121] O.G. Rossler, I. Bauer, H.Y. Chung, G. Thiel, Glutamate-induced cell death of immortalized murine hippocampal neurons: neuroprotective activity of heme oxygenase-1, heat shock protein 70, and sodium selenite, *Neurosci. Lett.* 362 (2004) 253–257.
- [122] H. Gutierrez, V.A. Hale, X. Dolcet, A. Davies, NF-kappaB signalling regulates the growth of neural processes in the developing PNS and CNS, *Development* 132 (2005) 1713–1726.

- [123] H.L. Pahl, Activators and target genes of Rel/NF-kappaB transcription factors, *Oncogene* 18 (1999) 6853–6866.
- [124] M.K. Meffert, D. Baltimore, Physiological functions for brain NF-kappaB, *Trends Neurosci.* 28 (2005) 37–43.
- [125] C.S. Simpson, B.J. Morris, Regulation of neuronal cell adhesion molecule expression by NF-kappa B, *J. Biol. Chem.* 275 (2000) 16879–16884.
- [126] J.L. Madrigal, M.A. Moro, I. Lizasoain, P. Lorenzo, A. Castrillo, L. Bosca, J.C. Leza, Inducible nitric oxide synthase expression in brain cortex after acute restraint stress is regulated by nuclear factor kappaB-mediated mechanisms, *J. Neurochem.* 76 (2001) 532–538.
- [127] M. Grilli, M. Ribola, A. Alberici, A. Valerio, M. Memo, P. Spano, Identification and characterization of a kappa B/Rel binding site in the regulatory region of the amyloid precursor protein gene, *J. Biol. Chem.* 270 (1995) 26774–26777.
- [128] J. Kraus, C. Borner, E. Giannini, V. Holtt, The role of nuclear factor kappaB in tumor necrosis factor-regulated transcription of the human mu-opioid receptor gene, *Mol. Pharmacol.* 64 (2003) 876–884.
- [129] B. Kaltschmidt, R.A. Linker, J. Deng, C. Kaltschmidt, Cyclooxygenase-2 is a neuronal target gene of NF-kB, *BMC Mol. Biol.* 3 (2002) 16.
- [130] C.A. Kassed, T.L. Butler, G.W. Patton, D.D. Demesquita, M.T. Navidomskis, S. Memet, A. Israel, K.R. Pennypacker, Injury-induced NF-kappaB activation in the hippocampus: implications for neuronal survival, *FASEB J.* 18 (2004) 723–724.
- [131] J. Schwamborn, A. Lindecke, M. Elvers, V. Horejschi, M. Kerick, M. Rafigh, J. Pfeiffer, M. Prullage, B. Kaltschmidt, C. Kaltschmidt, Microarray analysis of tumor necrosis factor alpha induced gene expression in U373 human glioblastoma cells, *BMC Genomics* 4 (2003) 46.
- [132] W. Zhang, I. Potrovita, V. Tarabin, O. Herrmann, V. Beer, F. Weih, A. Schneider, M. Schwaninger, Neuronal activation of NF-kappaB contributes to cell death in cerebral ischemia, *J. Cereb. Blood Flow Metab.* 25 (2005) 30–40.
- [133] K.J. Campbell, N.D. Perkins, Post-translational modification of RelA(p65) NF-kappaB, *Biochem. Soc. Trans.* 32 (2004) 1087–1089.
- [134] A. Routtenberg, It's about time, in: P.E. Gold, W.T. Greenough (Eds.), *Memory Consolidation*, American Psychological Association, 2000, pp. 17–34 (Chapter 2).
- [135] P.J. Meberg, W.R. Kinney, E.G. Valcourt, A. Routtenberg, Gene expression of the transcription factor NF-kB in hippocampus: regulation by synaptic activity, *Mol. Brain Res.* 38 (1996) 179–190.
- [136] M.K. Meffert, J.M. Chang, B.J. Wiltgen, M.S. Fanselow, D. Baltimore, NF-kappaB functions in synaptic signaling and behavior, *Nat. Neurosci.* 6 (10) (2003) 1072–1078.
- [137] B.C. Albensi, M.P. Mattson, Evidence for the involvement of TNF and NF-kappaB in hippocampal synaptic plasticity, *Synapse* 35 (2000) 151–159.
- [138] T. Suzuki, S. Mitake, K. Okumura-Noji, J.P. Yang, T. Fujii, T. Okamoto, Presence of NF-kB-like and I-kB-like immunoreactivities in postsynaptic densities, *NeuroReport* 8 (1997) 2931–2935.
- [139] B. Kaltschmidt, T. Deller, M. Frotscher, C. Kaltschmidt, Ultrastructural localization of activated NF-kappaB in granule cells of the rat fascia dentata, *NeuroReport* 11 (2000) 839–844.
- [140] R. Cantera, E. Roos, Y. Engstrom, Dif and cactus are colocalized in the larval nervous system of *Drosophila melanogaster*, *J. Neurobiol.* 38 (1999) 16–26.
- [141] R. Freudenthal, F. Locatelli, G. Hermitte, H. Maldonado, C. Lafourcade, A. Delorenzi, A. Romano, Kappa-B like DNA-binding activity is enhanced after spaced training that induces long-term memory in the crab *Chasmagnathus*, *Neurosci. Lett.* 242 (1998) 143–146.
- [142] M. Povelones, K. Tran, D. Thanos, R.T. Ambron, An NF-kappaB-like transcription factor in axoplasm is rapidly inactivated after nerve injury in Aplysia, *J. Neurosci.* 17 (1997) 4915–4920.
- [143] D. Sulejczak, M. Skup, Axoplasmic localisation of the NF kappa B p65 subunit in the rat brain, *Acta Neurobiol. Exp.* 60 (2000) 217.
- [144] M. Nonaka, X.H. Chen, J.E. Pierce, M.J. Leoni, T.K. McIntosh, J.A. Wolf, D.H. Smith, Prolonged activation of NF-kappaB following traumatic brain injury in rats, *J. Neurotrauma* 16 (1999) 1023–1034.
- [145] I. Ferrer, E. Marti, E. Lopez, A. Tortosa, NF-kB immunoreactivity is observed in association with beta A4 diffuse plaques in patients with Alzheimer's disease, *Neuropathol. Appl. Neurobiol.* 24 (1998) 271–277.
- [146] H. Wellmann, B. Kaltschmidt, C. Kaltschmidt, Retrograde transport of transcription factor NF-kB in living neurons, *J. Biol. Chem.* 276 (2001) 11821–11829.
- [147] R. Schmied, R.T. Ambron, A nuclear localization signal targets proteins to the retrograde transport system, thereby evading uptake into organelles in aplysia axons, *J. Neurobiol.* 33 (1997) 151–160.
- [148] S. Hanz, M. Fainzilber, Integration of retrograde axonal and nuclear transport mechanisms in neurons: implications for therapeutics, *Neuroscientist* 10 (2004) 404–408.
- [149] L. Guerrini, A. Molteni, T. Wirth, B. Kistler, F. Blasi, Glutamate-dependent activation of NF-kappaB during mouse cerebellum development, *J. Neurosci.* 17 (1997) 6057–6063.
- [150] R. Schmidt-Ullrich, S. Memet, A. Lilienbaum, J. Feuillard, M. Raphael, A. Israel, NF-kB activity in transgenic mice: developmental regulation and tissue specificity, *Development* 122 (1996) 2117–2128.
- [151] V. Fridmacher, B. Kaltschmidt, B. Goudeau, D. Ndiaye, F.M. Rossi, J. Pfeiffer, C. Kaltschmidt, A. Israel, S. Memet, Forebrain-specific neuronal inhibition of nuclear factor-kappaB activity leads to loss of neuroprotection, *J. Neurosci.* 23 (2003) 9403–9408.
- [152] A.L. Bhakar, L.L. Tannis, C. Zeindler, M.P. Russo, C. Jobin, D.S. Park, S. MacPherson, P.A. Barker, Constitutive nuclear factor-kappa B activity is required for central neuron survival, *J. Neurosci.* 22 (2002) 8466–8475.
- [153] S.D. Speese, N. Trotta, C.K. Rodesch, B. Aravamudan, K. Broadie, The ubiquitin proteasome system acutely regulates presynaptic protein turnover and synaptic efficacy, *Curr. Biol.* 13 (2003) 899–910.
- [154] I. Ninan, O. Arancio, Presynaptic CaMKII is necessary for synaptic plasticity in cultured hippocampal neurons, *Neuron* 42 (2004) 129–141.
- [155] Y. Ouyang, D. Kantor, K.M. Harris, E.M. Schuman, M.B. Kennedy, Visualization of the distribution of autophosphorylated calcium/calmodulin-dependent protein kinase II after tetanic stimulation in the CA1 area of the hippocampus, *J. Neurosci.* 17 (1997) 5416–5417.
- [156] M.N. Scholzke, I. Potrovita, S. Subramaniam, S. Prinz, M. Schwaninger, Glutamate activates NF-kappaB through calpain in neurons, *Eur. J. Neurosci.* 18 (2003) 3305–3310.
- [157] Y. Ben-Neriah, M.L. Schmitz, Of mice and men, *EMBO Rep.* 5 (2004) 668–673.
- [158] M.P. Mattson, Y. Goodman, H. Luo, W. Fu, K. Furukawa, Activation of NF-kappaB protects hippocampal neurons against oxidative stress-induced apoptosis: evidence for induction of manganese superoxide dismutase and suppression of peroxynitrite production and protein tyrosine nitration, *J. Neurosci. Res.* 49 (1997) 681–697.
- [159] P. Fernyhough, D.R. Smith, J. Schapansky, R. Van Der Ploeg, N.J. Gardiner, C.W. Tweed, A. Kontos, L. Freeman, T.D. Purves-Tyson, G.W. Glazner, Activation of nuclear factor-kappaB via endogenous tumor necrosis factor {alpha} regulates survival of axotomized adult sensory neurons, *J. Neurosci.* 25 (2005) 1682–1690.
- [160] C.L. Masters, G. Simms, N.A. Weinman, G. Multhaup, B.L. McDonald, K. Beyreuther, Amyloid plaque core protein in Alzheimer disease and Down syndrome, *Proc. Natl. Acad. Sci. U. S. A.* 82 (1985) 4245–4249.
- [161] J. Kang, H.G. Lemaire, A. Unterbeck, J.M. Salbaum, C.L. Masters, K.H. Grzeschik, G. Multhaup, K. Beyreuther, H.B. Muller, The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor, *Nature* 325 (1987) 733–736.
- [162] J. Hardy, D.J. Selkoe, The amyloid hypothesis of Alzheimer's

- disease: progress and problems on the road to therapeutics, *Science* 297 (2002) 353–356.
- [163] M.P. Mattson, Pathways towards and away from Alzheimer's disease, *Nature* 430 (2004) 631–639.
- [164] A. Ravati, B. Ahlemeyer, A. Becker, S. Klumpp, J. Kriegstein, Preconditioning-induced neuroprotection is mediated by reactive oxygen species and activation of the transcription factor nuclear factor-kappaB, *J. Neurochem.* 78 (2001) 909–919.
- [165] N. Blondeau, C. Widmann, M. Lazdunski, C. Heurteaux, Activation of the nuclear factor-kappaB is a key event in brain tolerance, *J. Neurosci.* 21 (2001) 4668–4677.
- [166] D. Baltimore, Intracellular immunization, *Nature* 335 (1988) 395–396.
- [167] B. Kaltschmidt, C. Kaltschmidt, T.G. Hofmann, S.P. Hehner, W. Droge, M.L. Schmitz, The pro- or anti-apoptotic function of NF-kappaB is determined by the nature of the apoptotic stimulus, *Eur. J. Biochem.* 267 (2000) 3828–3835.
- [168] H. Aleyasin, S.P. Cregan, G. Iyirhiaro, M.J. O'Hare, S.M. Callaghan, R.S. Slack, D.S. Park, Nuclear factor-(kappa)B modulates the p53 response in neurons exposed to DNA damage, *J. Neurosci.* 24 (2004) 2963–2973.
- [169] S. Hunot, B. Brugg, D. Ricard, P.P. Michel, M.P. Muriel, M. Ruberg, B.A. Faucheux, Y. Agid, E.C. Hirsch, Nuclear translocation of NF-kappaB is increased in dopaminergic neurons of patients with Parkinson disease, *Proc. Natl. Acad. Sci. U. S. A.* 94 (1997) 7531–7536.
- [170] C. Culmsee, J. Siewe, V. Junker, M. Retiounskaia, S. Schwarz, S. Camandola, S. El-Metainy, H. Behnke, M.P. Mattson, J. Kriegstein, Reciprocal inhibition of p53 and nuclear factor-kappaB transcriptional activities determines cell survival or death in neurons, *J. Neurosci.* 23 (2003) 8586–8595.
- [171] G.E. Hardingham, H. Bading, The Yin and yang of NMDA receptor signalling, *Trends Neurosci.* 26 (2003) 81–89.
- [172] B. Kaltschmidt, D. Ndiaye, M. Korte, S. Pothion, L. Laurence Arbibe, M. Prüllage, J. Pfeiffer, A. Lindecke, V. Staiger, A. Israël, C. Kaltschmidt, S. Memet, Essential role of the transcription factor NF-kappaB in memory and synaptic plasticity (submitted for publication).